

DERWENT-ACC-NO: 1996-045316
DERWENT-WEEK: 199605
COPYRIGHT 1999 DERWENT INFORMATION LTD

TITLE: Inhibitor of degradation of muco-polysaccharide, active oxygen inactivator and cosmetics - comprise super-oxide dismutase-like active oxygen removers obtd. from extract of flower petals

PATENT-ASSIGNEE: NARISU KESHOHIN KK[NARIN]

PRIORITY-DATA: 1994JP-0131057 (May 20, 1994)

PATENT-FAMILY:

PUB-NO	PUB-DATE	LANGUAGE	
PAGES	MAIN-IPC		
JP 07309770 A	November 28, 1995	N/A	011
A61K 035/78			

APPLICATION-DATA:

PUB-NO	APPL-DESCRIPTOR	APPL-NO
APPL-DATE		
JP07309770A	N/A	1994JP-0131057
May 20, 1994		

INT-CL (IPC): A61K007/00; A61K007/06 ; A61K007/48 ;
A61K035/78

ABSTRACTED-PUB-NO: JP07309770A

BASIC-ABSTRACT: Superoxide dismutase (SOD)-like active oxygen removers contg. at least one of extract of petals of flowers of 24 kinds of plants, and inhibitors of degradation of mucopolysaccharide contg. at least one of extract of petals of flowers of 42 kinds of plants including the preceding 24 kinds of plants. Cosmetics contg. at least one of the extracts of SOD-like active oxygen removers and inhibitors of degradation of mucopolysaccharide.

Petals of flowers of 42 plants (e.g. rose, peach, Japanese apricot, Thunberg spirea, sasanqua camellia, common camellia, torch azalea, kobus magnolia, yulan magnolia, Chinese paeony, carnation, snapdragon, daisy, dandelion, Japanese

wisteria, Chinese cabbage, common stock, hollyhock, shrub althea, cotton-rose hibiscus, common hydrangea, common crape myrtle and sweet-scented oleander) are extracted with water and/or lower alcohols (e.g. MeOH, EtOH and PrOH) and the extract is added in various bases (e.g. soln., emulsion, ointment, oil, wax, sol, gel and powder) including cosmetics and external preps..

USE/ADVANTAGE - Inhibitors of degradation of mucopolysaccharide, active oxygen removers and cosmetics. Prevention of ageing due to degradation of mucopolysaccharides with active oxygen and UV ray.

In an example, extracts of petals of flowers were tested for the inhibition of degradation of hyaluronic acid in ascorbic acid-Fe and H2O2-Fe systems at 10 and 0.1 mg/ml, respectively. Extract of pink rose showed inhibitory rate of 86.1 and 37.2%, respectively. Crape myrtle showed corresp. rate of 89.6 and 38.1 %, respectively.

CHOSEN-DRAWING: Dwg.0/0

TITLE-TERMS:

INHIBIT DEGRADE MUCO POLYSACCHARIDE ACTIVE OXYGEN INACTIVATE
COSMETIC COMPRISE
SUPER OXIDE DISMUTASE ACTIVE OXYGEN REMOVE OBTAIN EXTRACT FLOWER
PETAL

DERWENT-CLASS: B04 D21

CPI-CODES: B04-A08C2; B04-A10; B14-D09; B14-R01; B14-S08; D08-B;

CHEMICAL-CODES:

Chemical Indexing M1 *01*

Fragmentation Code

M423 M903 Q254 Q262 Q624 V400 V404 V802 V811

SECONDARY-ACC-NO:

CPI Secondary Accession Numbers: C1996-015009